



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

**STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES
OF THE UNITED STATES (Untabulated).**

CONNECTICUT.—Month of February, 1912. Population of reporting towns, 1,126,841. Total number of deaths from all causes 1,550, including diphtheria 9, measles 16, scarlet fever 8, tuberculosis (pulmonary) 118, typhoid fever 4. Cases reported: Diphtheria 143 in 33 towns, measles 781 in 50 towns, scarlet fever 194 in 45 towns, smallpox 1, tuberculosis (pulmonary) 168 in 38 towns, typhoid fever 28 in 18 towns.

FLORIDA.—Week ended March 2, 1912. Reports from the State board of health show diphtheria present in 4 localities with 10 cases, malaria in 2 localities with 8 cases, smallpox in 9 counties with 68 cases, tuberculosis in 9 localities with 15 cases, typhoid fever in 5 localities with 14 cases.

KANSAS.—Month of January, 1912. Population, 1,690,949. Total number of deaths from all causes not reported. The deaths include diphtheria 7, scarlet fever 6, smallpox 1, tuberculosis 21, typhoid fever 5. Cases reported: Diphtheria 91, measles 78, scarlet fever 202, smallpox 49, tuberculosis 209, typhoid fever 39.

PENNSYLVANIA.—Reports received from the State department of health show as follows:

Month of December, 1911. *Mortality.*—The total number of deaths was 9,307, including typhoid fever 142, scarlet fever 49, diphtheria 252, measles 29, whooping cough 64, smallpox 1, influenza 100, malaria 6, tuberculosis of the lungs 774, tuberculosis of other organs 118, cancer 439, diabetes 73, pellagra 1, meningitis 48, acute anterior poliomyelitis 6, pneumonia 1,253, diarrhea and enteritis (under 2 years) 196, diarrhea and enteritis (over 2 years) 65, Bright's disease 611, early infancy 520, suicide 61, accidents in mines 104, railway injuries 136, other forms of violence 462, all other diseases 3,797.

Month of January, 1912. *Morbidity.*—The total number of cases of communicable diseases reported was 13,462, including anterior poliomyelitis 11, anthrax 1, cerebrospinal meningitis 11, chickenpox 1,427, diphtheria 1,436, erysipelas 207, German measles 45, malarial fever 3, measles 4,242, mumps 688, pellagra 3, pneumonia 883, puerperal fever 11, rabies 1, scarlet fever 1,185, smallpox 15, tetanus 5, trachoma 5, trichiniasis 3, tuberculosis 1,329, typhoid fever 772, uncinariasis 4, whooping cough 1,175.

WISCONSIN.—Three months ended December 31, 1911. Population, 2,333,860. Total number of deaths from all causes 6,010, including diphtheria 93, measles 17, scarlet fever 14, tuberculosis 481, typhoid fever 93. Cases reported: Diphtheria 824, measles 856, scarlet fever 1,114, smallpox 231, tuberculosis (pulmonary) 236, typhoid fever 374.